

CONFIDENCE  
STARTS  
HERE

# Pylex<sup>®</sup>

Herbicide

## Strikingly Effective

- New and unique chemistry use pattern for Bentgrass greens
- Unmatched control of warm season grass and broadleaf weeds in cool season turf
- Targeted treatment with clear visual symptoms for exceptional performance and efficacy
- Highly selective herbicide presented in pre-measured syringes for easy application and greater turf safety

For more information on Pylex<sup>®</sup> Herbicide, visit [turf-solutions.basf.com.au](http://turf-solutions.basf.com.au) or contact your local BASF representative on **1800 558 399**

**■ - BASF**  
We create chemistry

# Pylex®

## Herbicide

Pylex Herbicide revolutionises the control of warm season grass and broadleaf weeds in cool season turf. This new and unique use pattern enables the safe and efficient removal of broadleaf weed species and warm season turf species, including Buffalo grass, Kikuyu, Couch and Queensland blue couch, in cool season turf on Bentgrass greens. As the only solution for selective weed control on greens, Pylex is pioneering an unmatched combination of turf safety and exceptional efficacy. Featuring pre-measured packaging and highly selective control, Pylex increases labour efficiencies and turf safety, minimising chemical calculation risks.

| Situation   | Weed Controlled   | Rate  | Critical Comments   |
|---|---|---|---|
| Bentgrass Greens  | Warm season grass weeds including:<br>Buffalo ( <i>Stenotaphrum secundatum</i> ),<br>Couch ( <i>Cynodon dactylon</i> ),<br>Hybrid couch ( <i>Cynodon dactylon</i> x <i>Cynodon transvaalensis</i> ),<br>Crowsfoot grass ( <i>Eleusine indica</i> ),<br>Kikuyu ( <i>Pennisetum clandestinum</i> ),<br>Qld blue couch / Summer grass ( <i>Digitaria</i> spp),<br>Broadleaf weeds including:                                 | 0.375 mL/100 m <sup>2</sup><br>in 4-6 L water<br>+ 0.5% MSO | Apply to actively growing weeds. Two repeat applications 21-28 days apart are required for optimum control. Some minor transient bleaching of Bentgrass may be observed 7-14 days post application.<br><br>For best results do not water or irrigate for 24 hours post application.   |
| Fine fescue, Tall fescue, Perennial Ryegrass, Kentucky blue grass | <i>Amaranthus</i> spp,<br>Bindy eye ( <i>Soliva sessilis</i> ),<br>Creeping oxalis ( <i>Oxalis corniculata</i> ),<br>Everlasting (cudweed) ( <i>Gamochaeta pensylvanica</i> ),<br>Fleabane/ Horseweed ( <i>Conyza bonariensis</i> ),<br>Scarlet Pimpernel ( <i>Anagallis arvensis</i> ),<br>Sowthistle ( <i>Sonchus oleraceus</i> ),<br>White clover ( <i>Trifolium repens</i> ),<br>Wild Carrot ( <i>Daucus carota</i> ) | 0.75 mL/100 m <sup>2</sup><br>in 4-6 L water<br>+ 0.5% MSO  | Apply to actively growing weeds.<br>Under some conditions a single application of Pylex may not be sufficient for complete control. A sequential application 21-28 days post application may be required.<br><br>Some minor transient bleaching of turf may be observed 7-14 days post application.<br><br>For best results do not water or irrigate for 24 hours post application. |

Refer to the product label for full directions for use table.

## Packaging

Available in 5x 0.375mL syringe packs.



For more information on Pylex® Herbicide, visit [turf-solutions.basf.com.au](https://turf-solutions.basf.com.au) or contact your local BASF representative on **1800 558 399**

**ALWAYS READ AND FOLLOW LABEL DIRECTIONS BEFORE USING ANY PRODUCT IN THIS FACT SHEET.**

This fact sheet is intended as general advice. Disclaimer: The information submitted in this publication is based on current BASF knowledge and experience. In view of the many factors that may affect its application, this data does not relieve the user from carrying out their own tests. The data does not imply assurance of certain properties or of suitability for a specific purpose. It is the responsibility of the user to ensure that any proprietary rights and existing laws and legislation are observed.