

POISON

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

STORM[®] Secure

Wax Block Rodenticide

ACTIVE CONSTITUENT: 0.05 g/kg FLOCOUMAFEN

Ready to use bait- kills in a single feed.

For control of mice in industrial, commercial, agricultural and domestic buildings and for control of rats in and around industrial, commercial, agricultural and domestic buildings.

Controls rats and mice resistant to warfarin.

IMPORTANT: READ THE SAFETY DIRECTIONS BEFORE USING THIS PRODUCT

CONTENTS: 500 g – 10 kg

BASF Australia Ltd
ABN 62 008 437 867
Level 23, 40 City Road Southbank VICTORIA 3006
pest-control.basf.com.au

® Registered trademark of BASF

APVMA Approval No.: 54191/RV2026

RESTRAINTS

DO NOT use in areas accessible to non-target animals, pets or children unless secured in a tamper resistant bait station. If bait can be dislodged from bait station, bait station must be secured in place.

DO NOT place bait outdoors unless it is in a tamper-resistant and weather resistant bait station which is secured in place.

DO NOT place bait stations more than 2 meters from buildings.

DO NOT place bait stations outdoors for control of mice.

DO NOT bait in areas where wildlife may be collected for human consumption.

DO NOT place in animal or livestock housing unless used in tamper-resistant and weather-resistant bait stations fixed to the ground or other structures.

DO NOT place bait or bait stations above areas which would allow them to contaminate pig and poultry food or drinking water.

DO NOT apply this product directly into burrows.

DIRECTIONS FOR USE

SITUATION	PEST	RATE	CRITICAL COMMENTS
In and around (within 2 m) industrial, commercial, agricultural and domestic buildings.	Rats	3 – 4 blocks per bait station	Baits must only be placed in areas frequented by the rats. Select suitable bait locations, such as along rat runs and under rubbish. Try to establish a barrier of bait locations between living and feeding areas. See baiting strategy instructions below.
In industrial, commercial, agricultural and domestic buildings.	Mice	1 – 2 blocks per bait station	Baits must only be placed in areas frequented by the mice. Place bait stations 2 metres apart in areas where activity is obvious, particularly where droppings are seen. Mice have very limited home ranges and do not need to drink. They are more difficult to control than rats because their feeding patterns are more erratic. Selection of bait locations is therefore even more important than for rats. See baiting strategy instructions below.

Baiting Strategy

Survey site prior to treatment, including assessment of the extent of infestation and identification of any indications of the presence of non-target animals.

Record the number and locations of bait stations for the purpose of inspection, replacement and later removal. Always use tamper-resistant bait stations that are secured in place in

child accessible areas and in animal or livestock housing. When using outdoors for control of rats, baits must be secured in a tamper-resistant and weather-resistant bait station which is secured in place. When used in food producing animal housing situations careful consideration is necessary regarding the placement of bait stations to ensure any rodenticide contamination of water, feed or livestock is prevented.

Inspect the bait locations after 3 days and select new locations where there has been no take. Inspect again 4 days later and replace eaten blocks. Repeat bait station inspections at 7-day intervals, replacing fresh blocks only when blocks have been eaten, until all signs of rodent activity have disappeared. Normally only 3 to 4 baiting rounds (pulses) are required.

DO NOT use the product beyond 35 days without an evaluation of the state of the infestation and of the efficacy of the treatment. Remove all uneaten bait at the end of treatment.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

GENERAL INSTRUCTIONS

Alternative food sources should be removed or reduced as much as possible. A lethal dose is readily consumed in a single feed but death may not occur for up to seven days.

PRECAUTIONS

Caution should be used when baiting near areas where hunting of wildlife may occur. Careful consideration of bait placement is necessary to avoid secondary exposure from contaminated wildlife that may be consumed.

DO NOT place bait in areas where there is a possibility of contaminating food, livestock feed or surfaces that come in direct contact with food or livestock feed.

PROTECTION OF LIVESTOCK AND WILDLIFE

Hazardous to livestock, poultry and wildlife. DO NOT place baits in locations that are accessible to domestic animals, livestock, non-target native animals or birds.

Search for and dispose of dead rodents and slugs/snails in the infested area at each visit to prevent secondary poisoning. In case slugs/snails are present, move bait station to another location within the rodent infested site, away from slugs/snails. Dispose of slugs/snails in a way non-target animals are not exposed. Dispose of dead rodents and uneaten bait in compliance with local, state or territory government regulations.

DO NOT allow livestock or poultry to consume dead or moribund mice and rats. All dead or moribund animals and rodent faeces from baited rodents MUST be removed from areas accessible to livestock or poultry.

DO NOT use this product to control native animals.

PROTECTION OF FISH, CRUSTACEANS AND ENVIRONMENT

Very toxic to aquatic life. DO NOT contaminate wetlands or watercourses with this product or used containers. Place the bait stations in areas not liable to flooding. When placing bait stations close to water drainage systems, ensure that bait contact with water is avoided.

STORAGE AND DISPOSAL

Store in the closed original container in a safe place. DO NOT store for prolonged periods in direct sunlight. Triple rinse containers before disposal. Break, crush or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available, dispose of empty packaging in compliance with relevant local, state or territory government regulations. Dispose of unused product in compliance with relevant local, state or territory government regulations. DO NOT burn empty containers or product.

SAFETY DIRECTIONS

Repeated minor exposure may have a cumulative poisoning effect. Do not touch bait, use scoop or measure. When using the product wear disposable gloves. If on skin and after each baiting, wash thoroughly with soap and water. During cleanup and disposal operations of bait and bait stations, wear a single layer of clothing and disposable gloves. Wear disposable gloves when handling rodent carcasses. Wash clothes after performing re-handling activities.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26, New Zealand 0800 764 766. Vitamin K1 (Phytomenadione) is antidotal.

CONDITIONS OF SALE

All conditions and warranties rights and remedies implied by law or arising in contract or tort whether due to the negligence of BASF Australia Ltd ABN 62008437867 or otherwise are hereby expressly excluded so far as the same may legally be done provided however that any rights of the Buyer pursuant to non excludable conditions or warranties of the Competition and Consumer Act 2010 or any relevant legislation of any State are expressly preserved but the liability of BASF Australia Ltd or any intermediate Seller pursuant thereto shall be limited if so permitted by the said legislation to the replacement of the goods sold or the supply of equivalent goods and all liability for indirect or consequential loss or damage of whatsoever nature is expressly excluded. This product must be used or applied strictly in accordance with the instructions appearing hereon. This product is solely sold for use in Australia and must not be exported without the prior written consent of BASF Australia Ltd.

APVMA Approval No: 54191/RV2026

Batch No:

Date of Manufacture:

® = Registered trademark of BASF

© BASF 2026

* Other registered trademarks

BASF Australia Ltd

ABN 62 008 437 867

Level 23, 40 City Road

Southbank VICTORIA 3006

pest-control.basf.com.au

FOR SPECIALIST ADVICE IN AN

EMERGENCY ONLY

PHONE 1800 803 440

TOLL FREE - ALL HOURS - AUSTRALIA WIDE

GHS STATEMENTS

May cause damage to organs (Blood) through prolonged or repeated exposure. If medical advice is needed, have product container or label at hand. Keep out of reach of children. Read carefully and follow all instructions. Do not breathe dust. Get medical advice/attention if you feel unwell.